



1/39

FIG. 1



Geldanamycin (3)



X=Cl Radical (1)

X=H Monocillin I (2)

FIG. 2





FIG. 3

3/39



- (a) TBDPSCI, imid., >95%; (b) DIBAL-H, -78 °C, 92%;
 (c) LiCl, DIPEA $(\text{EtO})_2\text{P}(\text{O})\text{CH}_2\text{CO}_2\text{Et}$, 95%;
 (d) DIBAL-H, -20 °C, 96%; (e) (+)-DET, $\text{Ti}(\text{O}i\text{Pr})_4$, TBHP, 90%, >95% ee; (f) $\text{SO}_3^*\text{pyridine}$, Et_3N , DMSO, 90%;
 (g) $\text{PH}_3\text{PCH}_3\text{Br}$, NaHMDS, 0 °C, 82%; (h) TBAF, 89%.




FIG. 4




FIG. 5





FIG. 6



a. *n*-BuLi, -78°C , 50% (6:1); b. TBSCl, 83%; c. H_2 C, 70%; d. (i) mCPBA, (ii) Ac_2O , Et_3N , H_2O , 60°C , (iii) NaHCO_3 , MeOH, 60%; e. SO_2Cl_2 , 50%

FIG. 7




FIG. 8




FIG. 9

9/39






FIG. 10



a. TBSCl, pyridine; b. NIS or NBS, TsOH; c. Pd(PPh)₃, RSnBu₃, d. nBu₄NF


FIG. 11-1




TO FIG. 11-2

FROM FIG. 11-1


FIG. 11-2




7




39




44




12/39



22




45




46

FIG. 12-1




FROM FIG. 12-1


FIG. 12-2



1. Rainey Ni





51




52

14/39




53



54

FIG. 13

15/39



- ^a (a) TBDPSCI, imid., >95%; (b) DIBAL-H, -78 °C, 92%; (c) LiCl, DIPEA (EtO)₂P(O)CH₂CO₂Et, 95%; (d) DIBAL-H -20 °C, 96%; (e) (+)-tetramethyltartaric acid diamide-BBu, Et₂Zn, CH₂I₂, 9 >95% ee; (f) SO₃*pyridine, Et₃N, DMSO, 90%; (g) Ph₃PCH NaHMDS, 0 °C, 82%; (h) TBAF, 89%; (i) 7, P(furyl)₃, DIA benzene, 60%

16/39

FIG. 14





FIG. 15-1



TO FIG. 15-2

FROM FIG. 15-1

FIG. 15-2






FIG. 16-1




19/39



TO FIG. 16-2

FROM FIG. 16-1


FIG. 16-2




21/39

FIG. 17-1


I. Radical




III. Cyclopropyl radical



II. Monocillin I



IV. Cyclopropyl monocillin



TO FIG. 17-2

FROM FIG. 17-1

FIG.17-2

MCF7 Cells Treated with Radicicol and Analogues




TO FIG. 17-3

23/39


FROM FIG. 17-2

FIG. 17-3


I. Radicol




VII. Radicol Oxime



V. Dimethyl Monocillin I




VI. Dimethyl Radicol




24/39

FIG. 18-1


I. Radical




III. Cyclopropyl radical



II. Monocillin I



IV. Cyclopropyl monocillin




TO FIG. 18-2

FROM FIG. 18-1


FIG.18-2

BT474 Cells Treated with Novel Radicicoliols (24hrs.)




HER2

FIG. 19




27/39

FIG. 20



28/39

FIG. 21



29/39

Therapeutic effect of Cycloproparadicol in nude mice bearing human mammary carcinoma MX-1 xenograft (Q2Dx6, iv.injection)




Fig. 22

Body weight changes of nude mice bearing human mammary carcinoma MX-1 xenograft:
Treatment with Cycloproparadicol (Q2Dx6, iv.injection)




Fig 23

31/39

Therapeutic effect of Radicolol & Cycloproparadicol in nude mice bearing human colon carcinoma (HCT-116) xenograft (QDx7, 4hr. iv-infusion)




Fig. 24

Body weight changes of human colon carcinoma (HCT-116) xenograft bearing nude mice following treatment with Radicolol & Cycloproparadicolol (QDx7, 4hr. iv-infusion)




Fig. 25

MX-1 tumors
12 hrs following a 6 hr CIVI




Fig. 26

34/39


IC₅₀ of Growth Inhibition of Different Tumor Cell lines

Fig. 27

35/39

Degradation of HER2 by Cycloproparadicicol Analogues




Fig. 28

36/39









Fig. 29

Cytotoxic effect on CCRF-CEM cell growth by radicicol analogs^a.



Compound	Structure	Cell growth inhibition (IC ₅₀ in μ M) ^b		
		CCRF-CEM	CCRF-CEM/VBL ^c	CCRF-CEM/taxol ^c
Radicicol (Sigma)		0.055 [±] 0.03	0.099 [1.8x]	0.070 [1.3x]
Cyclopropyl 1		4.81	9.84 [2.0x]	7.74 [1.6x]
Cyclopropyl 2		2.34	4.89 [2.1x]	2.89 [1.2x]
Cyclopropyl 3		0.58 [±] 0.13	0.87 [1.5x]	0.53 [0.9x]
Cyclopropyl 4 (Cycloproparadicicol)		0.055 [±] 0.04	0.041 [0.75x]	0.070 [1.3x]

37/39

Fig. 30A

38/39

Cytotoxic effect on CCRF-CEM cell growth by radicicol analogs^a. (Cont'd)

Compound	Structure	Cell growth inhibition (IC ₅₀ in μ M) ^b		
		CCRF-CEM	CCRF-CEM/VBL ^c	CCRF-CEM/taxol ^c
DechloroCyclopropa- radicicol Lactam		>10	>10	ND
Cycloproparadicicol Lactam		>5	>5	ND

^a Compounds of radicicol and cycloproparadicicol stereoisomers.^b Cell growth inhibition was measured by XTT tetrazonium assay after 72-hour incubation for cell growth. (Chou et al., Proc. Natl. Acad. Sci. USA 95: 15798-15802, 1998). Five to eight concentrations for each drug were used. IC₅₀ values were determined from dose-effect curves by using a computer program CalcuSyn for Windows by Chou and Hayball (Biosoft, Cambridge, UK, 1997).^c CCRF-CEM/VBL and CCRF-CEM/taxol are the CCRF-CEM sublines that are 320-fold and 42-fold resistant to vinblastine and taxol, respectively. Number in brackets is the fold of resistance of each drug when comparing the IC₅₀ values with those of the parent cell line, CCRF-CEM. The results showed that radicicol and cycloproparadicicol stereoisomers are not cross-resistant to vinblastine (typical MDR-Pgp substrate) nor to Taxol.

Fig. 30B

39/39




Figure 31: Her2 Degradation Assay

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record.**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.